



## Baltic Marine Environment Protection Commission

Tenth Meeting of the Seventh Baltic Sea Pollution Load  
Compilation (PLC-7) Project Implementation Group

PLC-7 IG 10-2020

Gothenburg, Sweden, 10-12 March 2020

---

<b>Document title</b>	Evaluation of the potential reduction of nutrient input from point sources
<b>Code</b>	4-2
<b>Category</b>	DEC
<b>Agenda Item</b>	4 - Current activities of the PLC-7 project and coordination with other HELCOM activities including ACTION project
<b>Submission date</b>	6.3.2020
<b>Submitted by</b>	Finland
<b>Reference</b>	

---

### Background

A draft assessment of effectiveness of measures to reduce nutrient loads from point sources was presented by SYKE, Finland at the PLC-7 9-2019. The Meeting invited national data reporters to verify the data in the report and provide feedback by 15 January 2020.

### Action requested

The Meeting is invited to consider the updated draft of the assessment report and agree on its presentation to PRESSURE 12-2020 for endorsement and publication.

## WP 4 Input of nutrients: effectiveness of measures

### WP 4.1 Following up existing measures

#### c) Contributions from point sources

##### Aim of the study

*Differences in the flow-normalized annual mean concentrations of discharges from direct point sources will be analysed in order to estimate the relative efficiency of the treatment of water discharges across the region. Inland (indirect) point sources will be assessed in regions where previous HELCOM analyses (HELCOM 2018c) have shown that these are a significant contribution to the total load. This analysis could indicate differences in the potential for measures across the region and variations between similar water treatment facilities in different countries.*

To estimate the reduction capacity of municipal wastewater treatment plants (MWWTP) PLC-7 data was collected from the PLC-database. The data includes both treatment plants discharging wastewaters directly to marine waters and to inland plants. Total phosphorus (PTOT) and total nitrogen (NTOT) loads of individual plants were divided by flows to get concentrations, which were compared to the limit values of the HELCOM recommendation 28E/5 and the EU urban wastewater directive (Table 1). Since the HELCOM limit values are stricter for PTOT than the respective EU values, the comparison was done with HELCOM recommendation values. If calculated concentration was above the limit value, the difference in mg/l was calculated back to tons for the estimation of the remaining reduction potential. Also retention of nutrients in inland waters was taken into account to get the estimate of the actual reduction potential benefitting the Baltic Sea.

Table 1. Municipal wastewater treatment requirements according to HELCOM recommendation 28E/5 and the respective requirements of the EU urban wastewater directive. PE = population equivalent.

PE	HELCOM				EU			
	PTOT		NTOT		PTOT		NTOT	
	mg/l	%	mg/l	%	mg/l	%	mg/l	%
300-2000	2	70	35	30				
2001-10000	1	80	35	30				
10001-100000	0.5	90	15	70-80	2	80	15	70-80
> 100000	0.5	90	10	70-80	1	80	10	70-80

## Material and Methods

### Data sources and calculation methods

Data of MWWTPs for the year 2017 (PLC-7 data) was collected from the PLC-database. Russia has only aggregated data, and also the Swedish data concerning inland MWWTPs was aggregated, but for this study Sweden submitted data of individual plants. The data consisted of altogether 3990 plants (Table 2).

Table 2. Number of MWWTPs in the PLC-database in 2017 by countries and sub-regions.

COUNTRY	BAP	WEB	KAT	SOU	GUF	GUR	ARC	BOB	BOS	Sum
DE	52	64								116
DK	44	319	217	42						622
EE					44	13				57
FI					162		36	113	83	394
LT	556						75			631
LV	9				1	28				38
PL	1685									1685
RU	4				16					20
SE	163		120	14				29	101	427
<b>Sum</b>	<b>2513</b>	<b>383</b>	<b>337</b>	<b>56</b>	<b>223</b>	<b>116</b>	<b>36</b>	<b>142</b>	<b>184</b>	<b>3990</b>

Population equivalent numbers (PE) were mostly missing in the database, but some countries (Denmark, Finland, Germany, Poland and Sweden) could submit this information enabling the classification of plants according PE numbers (Table 1). Since there is a high correlation between wastewater flow and PE ( $r^2$  0.81,  $n$  = 1741) in 2017, we used flow to estimate missing PE values according to this formula:

$$PE = \text{flow (m}^3/\text{a)} * 0.00904 + 4265$$

To be able to estimate the actual loads reaching the Baltic Sea retention in inland surface waters has to be taken into account. For that purpose MWWTPs were first divided into two categories: direct (zero retention) and indirect (variable retention depending on e.g. distance from sea and lake area). Since there is no estimate of retention of individual plants in the PLC-database other ways of retention were applied: A) for Danish plants 25% NTOT retention and 10% PTOT retention were used (Lars Svendsen personal communication). B) To estimate the retention for other countries MWWTP loads per sub-catchments were summed and the sums were compared with source apportionment figures (MWWTP loads reaching the Baltic Sea) derived from the PLC-7 data. C) Many countries (LT, LV, PL, RU) were lacking MWWTP loads in their source apportionment figures and for those countries published retention estimates were applied (Stålnacke et al. 2015, and Stålnacke Excel spreadsheet with P retention coefficients).

## Results

In 2017 the reported sum of MWWTP NTOT load in the PLC-database load was 69 800 t including both indirect and direct loads without retention taken into account (Table 3). Over half of the total municipal phosphorus loads originated from Poland (Table 4), which also has the biggest population. The respective nitrogen loads were divided more equally between the countries: Poland 28%, Russia 23% and Sweden 23%.

Loads were divided by country-wise population numbers to get concentrations (mg/l per inhabitants) enabling comparison of waste water treatment, or alternatively comprehensiveness of reporting of individual countries (Table 3). Based on this comparison Russia, Finland and Sweden had the highest per capita nitrogen concentrations. Denmark is known for its efficient nitrogen removal, which is reflected in low per capita nitrogen concentrations in outflowing wastewaters. Estonia, Latvia, Lithuania and Poland had nearly as low per capita nitrogen concentrations as Denmark, which may indicate that treatment is at a high level or part of the municipal nitrogen loads are missing from the PLC database.

Table 3. NTOT load of MWWTPs in 2017. **CHECK German population number!**

COUNTRY	BAP	WEB	KAT	SOU	GUF	GUR	ARC	BOB	BOS	Sum	%	Population	Population* connectivity	mg/inhabitant
	t	t	t	t	t	t	t	t	t	t				
DE	543	875								1419	2	4400000	4048000	350
DK	48	1065	850	1047						3010	4	5100000	4335000	694
EE					659	87				746	1	1300000	1066000	700
FI					3932		550	2809	2275	9566	14	5500000	4510000	2121
LT	1823						163			1986	3	2900000	2320000	856
LV	119				20	1181				1320	2	2200000	1672000	789
PL	19903									19903	28	38500000	27720000	718
RU	525				15506					16031	23	8400000	6972000	2299
SE	6387		3457	1005				1393	3614	15855	23	9500000	8265000	1918
<b>Sum</b>	<b>29348</b>	<b>1940</b>	<b>4307</b>	<b>2052</b>	<b>20118</b>	<b>1431</b>	<b>550</b>	<b>4202</b>	<b>5889</b>	<b>69836</b>	<b>100</b>	<b>77800000</b>	<b>60908000</b>	

<sup>1)</sup> Russian figures are based on aggregated data.

In 2017 the reported sum of MWWTP PTOT load in the PLC-database load was around 4330 t including both indirect and direct loads without retention taken into account (Table 4).

Table 4. PTOT load of MWWTPs in 2017. **CHECK German population number!**

COUNTRY	BAP	WEB	KAT	SOU	GUF	GUR	ARC	BOB	BOS	Sum	%	Population	Population* connectivity	mg/inhabitant
	t	t	t	t	t	t	t	t	t	t				
DE	37	27								64	2	4400000	4048000	15
DK	5	108	77	115						305	7	5100000	4335000	60
EE					31	5				36	1	1300000	1066000	28
FI					74		9	23	19	124	3	5500000	4510000	23
LT	136						15			151	4	2900000	2320000	52
LV	13				3	141				157	4	2200000	1672000	71
PL	2180									2180	52	38500000	27720000	57
RU	20				943					963	23	8400000	6972000	115
SE	104		70	26				11	31	243	6	9500000	8265000	26
<b>Sum</b>	<b>2495</b>	<b>136</b>	<b>147</b>	<b>142</b>	<b>1051</b>	<b>160</b>	<b>8.6</b>	<b>33.6</b>	<b>50.4</b>	<b>4223</b>	<b>100</b>	<b>77800000</b>	<b>60908000</b>	

<sup>1)</sup> Russian figures are based on aggregated data.

If all MWWTPs would follow HELCOM recommendation 28E/5, NTOT loads discharged into inland waters or directly to the Baltic Sea would decrease by 13 600 t (Table 5) and the respective PTOT loads would decrease by 2050 t (Table 6). In Estonia and Sweden there is no potential for further PTOT reductions. Finland, Sweden and Estonia had the lowest per capita phosphorus loads and Russia the highest.

Table 5. NTOT reduction potential in MWWTPs at source.

COUNTRY	BAP	WEB	KAT	SOU	GUF	GUR	ARC	BOB	BOS	Sum	
	t	t	t	t	t	t	t	t	t		%
DE	26	78								103	0.7
DK		4	3	0						7	0.0
EE					5					5	0.0
FI					1326		89	1703	1297	4415	30.5
LT	119					0				119	0.8
LV	8				9	207				224	1.5
PL	1375									1375	9.5
RU <sup>1)</sup>					4638					4638	32.0
SE	377		83	104				618	1498	2680	18.5
<b>Sum</b>	<b>1904</b>	<b>82</b>	<b>86</b>	<b>105</b>	<b>5978</b>	<b>207</b>	<b>89</b>	<b>2322</b>	<b>2795</b>	<b>13567</b>	<b>94</b>

<sup>1)</sup> Russian figures are based on aggregated data.

Table 6. PTOT reduction potential in MWWTPs at source.

COUNTRY	BAP	WEB	KAT	SOU	GUF	GUR	ARC	BOB	BOS	Sum	
	t	t	t	t	t	t	t	t	t		%
DE	11	2.6								14	0.7
DK	1.0	21	14	29						65	3.2
EE					0.1	0.0				0	0.0
FI					3.3		0.2	2.3	3.2	9	0.4
LT	48					4.9				53	2.6
LV	7.5				2.3	81				91	4.4
PL	1250									1250	61.0
RU <sup>1)</sup>	0.8				573					573	28.0
SE	0.1								0	0	0.0
<b>Sum</b>	<b>1319</b>	<b>24</b>	<b>14</b>	<b>29</b>	<b>578</b>	<b>86</b>	<b>0</b>	<b>2</b>	<b>3</b>	<b>2055</b>	<b>100</b>

<sup>1)</sup> Russian figures are based on aggregated data.

The reduction potential of loads discharged into inland waters would not totally benefit the Baltic Sea, since part of the loads would be retained along the route towards the sea. Approximately 17% of the NTOT reduction potential is lost, because of retention and 42% of the respective PTOT reduction potential (Tables 7 and 8). Thus, NTOT load into the Baltic Sea would decrease by 10500 t and the respective PTOT load by 1210 t, if all MWWTPs would follow HELCOM recommendation 28E/5.

For TN 88 % of the reductions potential is from Russia, Finland and Sweden while 86 % of the TP reduction potential are from Poland and Russia.

Table 7. NTOT reduction potential in MWWTPs at sea (taking into account retention in inland surface waters).

COUNTRY	BAP	WEB	KAT	SOU	GUF	GUR	ARC	BOB	BOS	Sum	
	t	t	t	t	t	t	t	t	t		%
DE1)	26	78								103	1.0
DK		3	2	0						6	0.1
EE					2					2	0.0
FI					908		89	1526	811	3334	31.6
LT	84					0				84	0.8
LV	7				4	148				158	1.5
PL	951									951	9.0
RU <sup>1)</sup>					3575					3575	33.9
SE	83		49	104				653	1387	2277	21.6
<b>Sum</b>	<b>1150</b>	<b>81</b>	<b>51</b>	<b>105</b>	<b>4489</b>	<b>148</b>	<b>89</b>	<b>2179</b>	<b>2199</b>	<b>10490</b>	<b>99</b>

<sup>1)</sup> Due to calculation methods German retention was estimated as zero.

<sup>2)</sup> Russian figures are based on aggregated data.

Table 8. PTOT reduction potential in MWWTPs at sea.

COUNTRY	BAP	WEB	KAT	SOU	GUF	GUR	ARC	BOB	BOS	Sum	
	t	t	t	t	t	t	t	t	t		%
DE <sup>1)</sup>	11	2.6								14	1.1
DK	1.0	21	13	29						63	5.3
EE					0.0	0.0				0	0.0
FI					2.1		0.2	2.2	2.3	7	0.6
LT	26					2.1				28	2.4
LV	4.1				1.2	50				55	4.6
PL	550									550	46.1
RU <sup>1)</sup>	0.6				494					495	41.4
SE	0.1								0	0	0.0
<b>Sum</b>	<b>593</b>	<b>23</b>	<b>13</b>	<b>29</b>	<b>497</b>	<b>52</b>	<b>0</b>	<b>2</b>	<b>2</b>	<b>1212</b>	<b>102</b>

<sup>1)</sup> Due to calculation methods German retention was estimated as zero.

<sup>2)</sup> Russian figures are based on aggregated data.

## Conclusions and recommendations

Even if municipal waste water treatment has improved substantially during the last decades there is still remarkable nutrient reduction potential in municipal waste water nutrient loads: NTOT load into the Baltic Sea would decrease by 10500 t and the respective PTOT load by 1210 t, if all MWWTPs would follow HELCOM recommendation 28E/5. This would correspond to nearly 10% of the BSAP reduction targets. For TN 88 % of the reductions potential is from Russia, Finland and Sweden while 86 % of the TP reduction potential are from Poland and Russia.

The uncertainties in estimating reduction potential were partly connected to the comprehensiveness of national nutrient loads: low nitrogen concentrations in treated waste waters indicated that part of the loads were missing from the PLC database. Therefore, there still is a need for further checking of possible gaps in data coverage.

Russia submitted only aggregated data disabling precise calculations of PE numbers and retention in inland waters. For other countries also only average catchment wise retention values were applied. Better knowledge about location of the plants and retention dynamics

would give more reliable estimates of how much of the potential reductions would actually benefit the Baltic Sea. This would be especially important for Polish and Russian plants.

## References

Stålnacke, P. Pengerud, A., Vassiljev, A., Smedberg, E., Mörth, C.-M., Hägg, H. E., C. Humborg, C. and. Andersen H. E. 2015. Nitrogen surface water retention in the Baltic Sea drainage basin. *Hydrol. Earth Syst. Sci.*, 19, 981–996.